ΑD)		

Award Number: DAMD17-96-1-6212

TITLE: Training in Support of Research Project Entitled "Tumor Cell De-adhesion by Aberrant, Single Skubmit Integrins"

PRINCIPAL INVESTIGATOR: Yan Zhang, Ph.D.

CONTRACTING ORGANIZATION: The Burnham Institute

La Jolla, California 92037

REPORT DATE: July 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20000707 016

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching edating data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Hoadquarters Services, Directorate for information Departies and Reports, 1215 Jefferson Davis Highway, Suite 1204, Artington, VA 22202-4302, and to the Office of

Manager And Conges - aperior - reconduct - re	COL COT OTTO LOCAL TEDORNINGSON, DIG 20000			
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND DATES COVERED		
	July 1999	Annual (1 Jul 9	8 - 30 Jun	99)
4. TITLE AND SUBTITLE Training in Support of Research Project Entitled "Tumor Cell De-adhesion by Aberrant, Single Subunit Integrins"			5. FUNDING NUMBERS DAMD17-96-1-6212	
6. AUTHOR(S) Yan Zhang, Ph.D.				
7. PERFORMING ORGANIZATION NA Burnham Institute La Jolla, California 92037	ME(S) AND ADDRESS(ES)		8. PERFORMING REPORT NUI	G ORGANIZATION MBER
E-MAIL:				
9. Sponsoring / Monitoring Agency Name(s) and Address(es)		10. SPONSORING / MONITORING AGENCY REPORT NUMBER		
U.S. Army Medical Research and I Fort Detrick, Maryland 21702-501				
11. SUPPLEMENTARY NOTES				
12a DISTRIBUTION / AVAILABILITY Approved for public rel		limited		12b. DISTRIBUTION CODE

13, ABSTRACT (Maximum 200 Words)

My purpose is try to identify a plasma membrane protein fingerprint which can be used to distinguish invasive breast cancer cell lines from non-invasive breast cancer lines. The methods included those for purification of plasma membrane proteins, extraction of proteins using urea/thiourea mixture, a standard procedure for 2D PAGE, excision of polypeptide "spots" from gels, trypsin digestion, MALDI TOF mass spectrometry and database searching for protein identity are well-established for this project. Three breast cancer cell lines (MDA-MB435, MDA-MB231 and MCF7) with different metastatic potential have been grown and 2-D maps of plasma membrane proteins have been generated in the laboratory. Computer-assisted analysis of the 2D polypeptide pattern showed that 56% of the polypeptide spots were the same between MCF7 and MDA-MB231 (two non-metastatic Breast Cancer cell lines), only 42% of the polypeptide spots were the same between MCF7 and MDA-MB435 (metastatic vs non-metastatic Breast Cancer cell lines). This differential expression of proteins is being characterized in order to determine which proteins are important to the metastatic condition. A number of proteins from the 2D gels has been identified by MALDI TOF mass spectrometry. Recently, antibody library phage display, has been expanded for further identifying the plasma membrane proteins.

14. SUBJECT TERMS Breast Cancer	15. NUMBER OF PAGES		
			16, PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	Unlimited

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18 298-102

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

X Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

X In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

N/A For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

PI - Signature

TABLE OF CONTENTS

FRONT COVER	1
REPORT DOCUMENTATION PAGE	2
FOREWORD	3
TABLE OF CONTENTS	4
PEPORT	5

Accomplishments towards Objectives

During the six months that I have worked on this project, progress has been made in numerous areas. With my assistance, many of the proteomics methodologies underwent final optimization and standard operating procedures were adopted. The methods included those for purification of plasma membrane proteins using cationic colloidal silica, extraction of proteins using urea/thiourea mixture and 2D PAGE parameters. In addition, a new method was established for spotting of matrix and peptides onto MALDI targets. This new technique, with its increased sensitivity and peptide yield, greatly advances our ability to identify proteins from 2D gels.

Specific Aim 1

Task 1. This task has not yet been completed. Although we have 3 adherent breast cancer cell lines in-house, we need to obtain the cell line for growth in suspension. The method for purification of plasma membranes using colloidal silica and extraction of the plasma membrane proteins using urea/thiourea has already been well-established in this lab.

Task 2. Since we do not yet have a breast cancer cell line for growth in suspension this task has not been addressed; however, a standard procedure for 2D PAGE of plasma membrane proteins has been adopted and the gel analysis software has been used extensively by the lab.

Specific Aim 2

Task 1. Although this task has not been accomplished, the techniques for excision of polypeptide "spots" from gels, trypsin digestion, MALDI TOF mass spectrometry and database searching for protein identity are well-established in the lab.

Specific Aim 3

Task 1. Three breast cancer cell lines (MDA-MB435, MDA-MB231 and MCF7) with different metastatic potential have been grown in the laboratory. More specifically, MCF7 is representative of a non-invasive mammary carcinoma as the cells retain epithelial morphology, require estrogen for growth and also fail to metastasize when implanted into the mammary fat pad of mice. In contrast, the MDA-MB435 cells exhibit a mesenchymal morphology, grow independently of estrogen and metastasize extensively from the mammary fat pad of mice. The third mammary carcinoma cell line, MDA-MB231, is tumorgenic but non-metastatic.

Task 2. Two-dimensional maps of plasma membrane proteins have been generated from the three breast cancer cell lines. On average, 376 polypeptide spots were resolved on each of the 2D gels. Computer-assisted analysis of the plasma membrane protein 2D polypeptide pattern showed that 56% of the polypeptide spots were the same between MCF7 and MDA-MB231 (two non-metastatic Breast Cancer cell lines), whereas only

Principal Investigator: Zhang, Yan

42% of the polypeptide spots were the same between MCF7 and MDA-MB435 (metastatic vs non-metastatic Breast Cancer cell lines). This differential expression of proteins at the plasma membrane is being characterized in order to determine which proteins are important to the metastatic condition.

Specific Aim 4

Task 1. This task has been initiated in that MALDI TOF mass spectrometry has been used to identify a number of proteins from the 2D gels of the plasma membrane proteins of metastatic and non-metastatic breast cancer cells.

Note: Specific Aim 4 has recently been expanded to include an additional methodology, antibody library phage display, for identifying the plasma membrane proteins that differ in expression between metastatic and non-metastatic breast cancer cells. This unique methodology will complement and further advance our ability to identify the complex array of plasma membrane proteins that contribute to the metastatic phenotype. In this approach RNA is extracted from the spleens of mice that have been immunized with the purified plasma membrane proteins of a cell line of interest. A cDNA library is then constructed by reverse transcription of the extracted RNA and a mouse Immunoglobulin G scFv (single chain variable Fragment) library is generated from this cDNA library. Finally, a phage display scFv library is constructed. The antibodies that display on the surface of the phage are then screened to identify the specific antibodies that bind the plasma membrane proteins immobilized on the blots of 2D gels. Similarities and differences between metastatic and non-metastatic breast cancer cell lines can be observed by identifying the differential binding of phage to the blots. The phage display strategy will be a powerful asset for obtaining antibodies to multiple proteins on the blots of 2D gels. In combination with the identification of these proteins by mass spectrometry, it is likely that I will be able to identify and raise antibodies to dozens if not hundreds of plasma membrane proteins at one time.

I am currently working out the logistics of the "phage display" approach using an endothelial cell line with known cell-surface markers as a positive control situation. Mice have been immunized and the sera evaluated by Enzyme-linked immunosorbent assay (ELISA) and by Fluorescence-assisted cell sorting (FACS). In both cases, the immune sera showed a positive response towards known endothelial cell-surface markers, whereas pre-immune sera did not. RNA has been isolated from the spleens of the immunized mice, a panel of IgG scFv library has been generated and a phage display scFv antibody library is currently being constructed.